# Pelvic Floor Impact Questionnaire – Short Form 7

**Instructions:** Some women find that bladder, bowel, or vaginal symptoms affect their activities, relationships, and feelings. For each question place an **X** in the response that best describes how much your activities, relationships, or feelings have been affected by your bladder, bowel, or vaginal symptoms or conditions <u>over the last 3 months</u>. Please make sure you mark an answer in <u>all 3 columns</u> for each question.

con foll	w do symptoms or ditions relate to the owing $\rightarrow$ ally affect your $\downarrow$	Bladder or Urine	Bowel or rectum	Vagina or pelvis
1.	Ability to do household chores (cooking, housecleaning, laundry)?	Not at all Somewhat Moderately Ouite a bit	Not at all Somewhat Moderately Ouite a bit	Not at all Somewhat Moderately Ouite a bit
2.	Ability to do physical activities such as walking, swimming, or other exercise?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
3.	Entertainment activities such as going to a movie or concert?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
4.	Ability to travel by car or bus for a distance greater than 30 minutes away from home?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
5.	Participating in social activities outside your home?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
6.	Emotional health (nervousness, depression, etc.)?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit
7.	Feeling frustrated?	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit	Not at all Somewhat Moderately Quite a bit

# Scoring the PFIQ-7:

All of the items use the following response scale:

0, Not at all; 1, somewhat; 2, moderately; 3, quite a bit

# Scales:

Urinary Impact Questionnaire (UIQ-7): 7 items under column heading "Bladder or urine" Colorectal-Anal Impact Questionnaire (CRAIQ-7): 7 items under column heading "Bowel or rectum" Pelvic Organ Prolapse Impact Questionnaire (POPIQ-7): 7 items under column heading "Pelvis or vagina"

#### Scale scores:

Obtain the mean value for all of the answered items within the corresponding scale (possible value 0 to 3) and then multiply by (100/3) to obtain the scale score (range 0 to 100). Missing items are dealt with by using the mean from answered items only.

# **PFIQ-7 Summary Score:**

Add the scores from the 3 scales together to obtain the summary score (range 0 to 300).

Reprinted with permission from: Barber MD, Walters MD, Bump RC. Short forms of two specific quality-of-life questionnaires for women with pelvic floor disorders (PFDI-20 and PFIQ-7. *Am J Obstet Gynecol*. 2005; 193:103-113; posted with permission from Elsevier, http://www.ajog.org/article/S0002-9378(04)02108-8/abstract.